The background is a dark gray color with several light gray butterfly silhouettes scattered across it. One butterfly in the bottom right corner is a vibrant blue color, while all others are light gray.

# Measure Of Central Tendency

## Introduction:

- In statistics, a central tendency is a central value or a typical value for a probability distribution.
- It is occasionally called an average or just the center of the distribution.
- The most common measures of central tendency are the arithmetic mean, the median and the mode.
- Measures of central tendency are defined for a population (large set of objects of a similar nature) and for a sample (portion of the elements of a population).

## Some Definitions

**Simpson and Kafka** defined it as “A measure of central tendency is a typical value around which other figures gather”

**Waugh** has expressed “An average stand for the whole group of which it forms a part yet represents the whole”.

**In layman's term,** a measure of central tendency is an **AVERAGE**. It is a single number of value which can be considered typical in a set of data as a whole.

# Importance Of Central Tendency

- To find representative value
- To make more concise data
- To make comparisons
- Helpful in further statistical analysis

# Mean

- The MEAN of a set of values or measurements is the sum of all the measurements divided by the number of measurements in the set.
- The mean is the most popular and widely used. It is sometimes called the arithmetic mean.

## Mean for Ungrouped data

- If we get the mean of the sample, we call it the sample mean and it is denoted by (read “x bar”).

$$\text{Sample mean} = \frac{\text{Sum of all the values in the sample}}{\text{Number of values in the sample}}$$

- If we compute the mean of the population, we call it the parametric or population mean, denoted by  $\mu$  (read “mu”).

$$\text{Population mean} = \frac{\text{Sum of all the values in the population}}{\text{Number of values in the population}}$$

# Arithmetic Mean Calculated Methods for grouped data:

- **Direct Method :**

$$\bar{X} = \frac{\sum fm}{N}$$

- **Short cut method :**

$$\bar{X} = A + \frac{\sum fd}{N}$$

- **Step deviation Method :**

$$\bar{X} = A + \frac{\sum fd}{N} \times i$$

## Example Of A.M:

A sample of five executives received the following bonus last year (\$000):

14.0, 15.0, 17.0, 16.0, 15.0

**Solution:**

$$\bar{X} = \frac{\Sigma X}{n} = \frac{14.0 + \dots + 15.0}{5} = \frac{77}{5} = 15.4$$

# Weighted Mean

- **Weighted mean** is the mean of a set of values wherein each value or measurement has a different weight or degree of importance. The following is its formula:

$$\bar{x} = \frac{\sum xw}{\sum w}$$

where  $\bar{x}$  = mean

x = measurement or value

w = number of measurements

# Example Of W.M

Below are Amaya's subjects and the corresponding number of units and grades she got for the previous grading period. Compute her grade point average.

Subject	Units	Grade
Filipino	.9	86
English	1.5	85
Mathematics	1.5	88
Science	1.8	87
Social Studies	.9	86
TLE	1.2	83
MAPEH	1.2	87

$$\bar{X} = \frac{(0.9 \cdot 86) + (1.5 \cdot 85) + (1.5 \cdot 88) + (1.8 \cdot 87) + (0.9 \cdot 86) + (1.2 \cdot 83) + (1.2 \cdot 87)}{9} = 86.1$$

Amaya's average grade is 86.1

# Harmonic Mean

- Harmonic mean is quotient of “number of the given values” and “sum of the reciprocals of the given values”.
- For Ungrouped Data

$$H.M \text{ of } X = \bar{X} = \frac{n}{\sum \left( \frac{1}{x} \right)}$$

- For grouped Data

$$H.M \text{ of } X = \bar{X} = \frac{\sum f}{\sum \left( \frac{f}{x} \right)}$$

# Harmonic Mean Example

Calculate the harmonic mean of the numbers: 13.2, 14.2, 14.8, 15.2 and 16.1

**Solution:**

The harmonic mean is calculated as below:

**AS**

$$H.M \text{ of } X = \bar{X} = \frac{n}{\sum \left( \frac{1}{x} \right)}$$

$$H.M \text{ of } X = \bar{X} = \frac{5}{0.3417} = 14.63$$

<b>X</b>	<b><math>\frac{1}{X}</math></b>
13.2	0.0758
14.2	0.0704
14.8	0.0676
15.2	0.0658
16.1	0.0621
Total	$\sum \frac{1}{X} = 0.3147$

**Example: Calculate the harmonic mean for the given below:**

Marks	30-39	40-49	50-59	60-69	70-79	80-89	90-99
F	2	3	11	20	32	25	7

**Solution:** Now  
We'll find H.M as:

$$\bar{X} = \frac{\sum f}{\sum \left(\frac{f}{x}\right)} = \frac{100}{1.4368} = 69.60$$

Marks	x	f	$\frac{f}{x}$
30-39	34.5	2	0.0580
40-49	44.5	3	0.0674
50-59	54.5	11	0.2018
60-69	64.5	20	0.3101
70-79	74.5	32	0.4295
80-89	84.5	25	0.2959
90-99	94.5	7	0.0741
<b>Total</b>		<b>4.5</b>	<b>0.4295</b>

# Geometric Mean

- Geometric mean is a kind of average of a set of numbers that is different from the arithmetic average.
- The geometric mean is well defined only for sets of positive real numbers. This is calculated by multiplying all the numbers (call the number of numbers  $n$ ), and taking the  $n$ th root of the total.
- A common example of where the geometric mean is the correct choice is when averaging growth rates.
- The geometric mean is NOT the arithmetic mean and it is NOT a simple average.
- **Mathematical definition:** The  $n$ th root of the product of  $n$  numbers.

# Formulas

- For grouped data:

$$G = \text{Anti} \left( \sum \frac{\log x}{n} \right)$$

- For Grouped data:

$$G = \text{anti} \left( \sum \frac{f \log x}{n} \right)$$

Question 1: Find the geometric mean of the following values:

15, 12, 13, 19, 10

**Solution:**

Given data, 15, 12, 13, 19, 10

$$n=5, \sum \log x=5.648$$

$$\text{As } G = \text{Anti}\left(\frac{\sum \log x}{n}\right)$$

$$G = \text{Anti}\left(\frac{5.648}{5}\right)$$

$$G = \text{Anti}(1.129)$$

$$G = 13.48$$

$x$	$\text{Log } x$
15	1.1761
12	1.0792
13	1.1139
19	1.2788
10	1.0000
Total	5.648

**MEDIAN**



# Median

- The **MEDIAN**, denoted **Md**, is the middle value of the sample when the data are ranked in order according to size.
- **Connor has defined as** “ The median is that value of the variable which divides the group into two equal parts, one part comprising of all values greater, and the other, all values less than median”
- **For Ungrouped data median is calculated as:**

$$\text{Median}(M) = \text{Size of } \left(\frac{N+1}{2}\right) \text{th item}$$

- **For Grouped Data:**

$$\text{Median}(M) = L1 + \frac{\frac{N}{2} - cf}{f} \times i$$

# Example OF Median

The ages for a sample of five college students are:

21, 25, 19, 20, 22

**Solution:**

Arranging the data in ascending order gives:  
19, 20, 21, 22, 25.

Here  $n=5$

As median =  $\left(\frac{N+1}{2}\right)$ th value

$Md = \left(\frac{5+1}{2}\right)$ th value

$= \left(\frac{6}{2}\right)$ th value

$= 3^{\text{rd}}$  value

Thus the median is **21**.

## Example of median For Grouped Data

*Solution :*

$$\text{As median} = L + \frac{h}{f} \left( \frac{n}{2} - c \right)$$

$$n=50; h=5; L=14.5$$

$$f=17 \text{ and } c=24$$

so

$$\begin{aligned} \text{Median} &= 14.5 + \frac{5}{17} \left( \frac{50}{2} - 24 \right) \\ &= 14.94 \end{aligned}$$

Marks	No. of students	<u>Less than</u> CF
0 – 4	2	2
5 – 9	8	10
10 – 14	14	24
15 – 19	17	41
20 – 24	9	50
<b>Total</b>	<b>50</b>	

# Mode

- The MODE, denoted  $M_o$ , is the value which occurs most frequently in a set of measurements or values. In other words, it is the most popular value in a given set.
- Croxton and Cowden : defined it as “the mode of a distribution is the value at the point where the items tend to most heavily concentrated. It may be regarded as the most typical of a series of value”

**Example 2:** In a crash test, 11 cars were tested to determine what impact speed was required to obtain minimal bumper damage. Find the mode of the speeds given in miles per hour below.

24, 15, 18, 20, 18, 22, 24, 26, 18, 26, 24



**Solution:** Ordering the data from least to greatest, we get:

15, 18, 18, 18, 20, 22, 24, 24, 24, 26, 26

**Answer:** Since both 18 and 24 occur three times, the modes are 18 and 24 miles per hour.

# Formula:

For Grouped Data:

$$\text{Mode} = L + \left( \frac{f_n - f_1}{(f_n - f_1) + (f_n - f_2)} \right) * h$$

- L is the lower class limit of the modal class
- $f_n$  is the frequency of the modal class
- $f_1$  is the frequency of the class before the modal class in the frequency table
- $f_2$  is the frequency of the class after the modal class in the frequency table
- h is the class interval of the modal class

Find the modal class and the actual mode of the data set below

Number	Frequency
1 - 3	7
4 - 6	6
7 - 9	4
10 - 12	2
13 - 15	2
16 - 18	8
19 - 21	1
22 - 24	2
25 - 27	3
28 - 30	2

Solution:

Here modal class=10-12

**L = 10, fn = 9, f1 = 4, f2 = 2, h = 3**

$$\text{Mode} = L + \left( \frac{fn - f1}{(fn - f1) + (fn - f2)} \right) * h$$

$$= 10 + \left( \frac{9 - 4}{(9 - 4) + (9 - 2)} \right) * 3$$

$$= 10 + \left( \frac{5}{5 + 7} \right) * 3$$

$$= 10 + 1.25$$

$$\text{Mode} = 11.25$$

## Advantages of Mode :

- Mode is readily comprehensible and easily calculated
- It is the best representative of data
- It is not at all affected by extreme value.
- The value of mode can also be determined graphically.
- It is usually an actual value of an important part of the series.

## Disadvantages of Mode :

- It is not based on all observations.
- It is not capable of further mathematical manipulation.
- Mode is affected to a great extent by sampling fluctuations.
- Choice of grouping has great influence on the value of mode.

# Advantages of Median:

- Median can be calculated in all distributions.
- Median can be understood even by common people.
- Median can be ascertained even with the extreme items.
- It can be located graphically
- It is most useful dealing with qualitative data

## Disadvantages of Median:

- It is not based on all the values.
- It is not capable of further mathematical treatment.
- It is affected fluctuation of sampling.
- In case of even no. of values it may not the value from the data.

# Properties of mode

- It is used when you want to find the value which occurs most often.
- It is a quick approximation of the average.
- It is an inspection average.
- It is the most unreliable among the three measures of central tendency because its value is undefined in some observations.

# Properties of Mean

1. Mean can be calculated for any set of numerical data, so it always exists.
2. A set of numerical data has one and only one mean.
3. Mean is the most reliable measure of central tendency since it takes into account every item in the set of data.
4. It is greatly affected by extreme or deviant values (*outliers*)
5. It is used only if the data are interval or ratio.

# Relations Between the Measures of Central Tendency


- In symmetrical distributions, the median and mean are equal

For normal distributions, mean = median = mode

- In positively skewed distributions, the mean is greater than the median



- In negatively skewed distributions, the mean is smaller than the median



# Conclusion

- A measure of central tendency is a measure that tells us where the middle of a bunch of data lies.
- Mean is the most common measure of central tendency. It is simply the sum of the numbers divided by the number of numbers in a set of data. This is also known as average.
- Median is the number present in the middle when the numbers in a set of data are arranged in ascending or descending order. If the number of numbers in a data set is even, then the median is the mean of the two middle numbers.
- Mode is the value that occurs most frequently in a set of data.

*Thank You...* 😊 😊

